



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Reports of States and yearly and monthly reports of cities of the United States.

COLORADO—*Denver*.—Month of January, 1900. Estimated population, 170,000. Total number of deaths, 218, including diphtheria, 3; enteric fever, 1; measles, 7; scarlet fever, 2, and 55 from tuberculosis.

MARYLAND—*Baltimore*.—Month of December, 1899. Estimated population, 541,000—white, 463,000; colored, 78,000. Total number of deaths, 769—white, 598; colored, 191, including diphtheria, 50; enteric fever, 13; scarlet fever, 1, and 83 from phthisis pulmonalis.

MASSACHUSETTS—*Lawrence*.—Month of January, 1900. Estimated population, 59,072. Total number of deaths, 99, including enteric fever, 1; whooping cough, 1, and 10 from phthisis pulmonalis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended February 17, 1900, from 67 observers, indicate that intermittent fever, pleuritis, and pneumonia increased and scarlet fever decreased in area of prevalence. Phthisis pulmonalis was reported present at 173, measles at 91, scarlet fever at 73, enteric fever at 33, whooping cough at 27, diphtheria at 20, cerebro-spinal meningitis at 4, and smallpox at 3 places.

MINNESOTA—*Stillwater*.—Month of January, 1900. Estimated population, 14,000. Total number of deaths, 15, including enteric fever, 1, scarlet fever, 1, and 2 from phthisis pulmonalis.

NEW YORK—*Jamestown*.—Year ended March 31, 1899. Census population, 16,038. Total number of deaths, 298, including diphtheria, 4, enteric fever, 5, and 1 from scarlet fever.

Rochester.—Month of January, 1900. Estimated population, 180,000. Total number of deaths, 173, including diphtheria, 4; enteric fever, 4; scarlet fever, 2, and 19 from phthisis pulmonalis.

OHIO.—Reports to the State board of health for the five weeks ended December 2, 1899, from 68 localities, having an aggregate estimated population of 1,301,773, show 43 deaths from diphtheria, 40 from enteric fever, scarlet fever, 15, and 6 from whooping cough.

Cleveland.—Month of January, 1900. Estimated population, 395,000. Total number of deaths, 466, including diphtheria, 25; enteric fever, 7; scarlet fever, 2; smallpox, 1; whooping cough, 1, and 11 from phthisis pulmonalis.

PENNSYLVANIA—*Erie*.—Year ended December 31, 1899. Census population, 40,634. Total number of deaths, 730, including diphtheria, 16; enteric fever, 18; measles, 19; scarlet fever, 9; whooping cough, 3, and 80 from tuberculosis.

Oil City.—Month of January, 1900. Estimated population, 17,482. Total number of deaths, 17, including enteric fever 2, and 2 from tuberculosis.

VIRGINIA—*Richmond*.—Month of January, 1900. Estimated population, 101,560. Total number of deaths, 141, including diphtheria, 1; enteric fever, 2, and 3 from tuberculosis.

WASHINGTON—*Seattle*.—Month of January, 1900. Estimated population, 90,000. Total number of deaths, 62, including enteric fever, 4; scarlet fever, 1, and 4 from tuberculosis.

WISCONSIN—*La Crosse*.—Two weeks ended February 17, 1900. Estimated population, 30,000. Total number of deaths, 20. No deaths from contagious diseases.